

Feet & Pregnancy

As the Pelvis Loosens, So Do The Feet


Body Changes During Pregnancy

Pregnancy triggers a wide range of changes in a women's body. The additional weight in the torso has immediate effects such as increased curvature in the lower back and a forward tilt of the pelvis. When a women's center of gravity is altered, the weight bearing stance also changes. This can cause increased demand on the muscles of the hips, legs and ankles. These changes contribute to the characteristic posture and gait associated with the pregnant women¹. The fact that feet change during pregnancy comes as no surprise to most women. Even if they are pregnant for the first time, most women have heard stories or warnings about increasing shoe sizes, foot pain and swelling. So what are the facts behind these problems and can these problems be helped?

Foot Problems Associated With Pregnancy

A common complaint – often overlooked – of women during pregnancy is foot pain. Women adjust their pattern of walking during pregnancy to improve stability. This is done to accommodate for obvious shifts in weight distribution and center of gravity, and requires muscles in the hips, pelvis, legs and ankles to work differently.

There are two common changes associated with the foot during pregnancy. The first is an increase in volume or size caused by edema (swelling) and the second is an increase in size caused by a looseness or laxity of the ligaments that support your foot. This ligament laxity is caused by a hormone called relaxin which is the same hormone that is loosening the ligaments of the birth canal and pelvis in preparation for delivery. This hormone does not target only the birthing ligaments, and has wide ranging effects throughout the body. The ligaments that hold the 26 bones of your foot can be



Increased Low Back Curve

Knees: move closer together & backwards

During pregnancy, the feet flatten and spread out from side to side and length-wise

Hormonal changes during pregnancy which aid the necessary pelvic expansion during birth also relax ligaments in the feet. This plus weight gain can flatten arches, leading to changes at the knee, hip and back.

affected, and as these ligaments loosen the arch of the foot collapses or over-pronates. Since the levels of relaxin are elevated in early pregnancy, these changes can occur over a period of months.

Also common during pregnancy are painful calluses, corns and cracked heels. These complaints are indicative of the underlying problem of your foot flattening and spreading out while your shoes remain the same size.

Are These Changes Permanent?

Studies in the Journal of the American Podiatric Medical Association have shown that the changes in the way a women walks during pregnancy have lasting effects and may not return to nor-

Feet & Pregnancy

As the Pelvis Loosens, So Do The Feet

mal following birth². Some experts feel that the shoe size changes associated with pregnancy are permanent³.

How Is It treated?

Unfortunately for many women, problems associated with their feet are often not addressed adequately. Complaints in the feet, knees and hips are often deemed secondary and dismissed in the hope that the problem will resolve spontaneously after birth⁴. This may be because many clinicians are concerned that the treatment protocols for these problems in non-pregnant women are inappropriate and even dangerous in pregnant women. Unfortunately this can lead to a treatment strategy of benign neglect.

The first step in addressing your foot pain is to remove any potential interference with your changing foot. To help with the swelling, soak your feet in cold water and prop them up to help with drainage. Buy a few new pairs of roomy comfortable shoes. Be aware that your feet may continue to change with each pregnancy. Avoid the use of shoes with heels as these only accentuate the problems in your foot at a time when it is at its weakest structurally.

Custom Foot Orthotics

The over-pronation associated with pregnancy is best treated with a truly custom, biomechanically correct foot orthotic. In fact, pregnancy provides an opportunity to address foot problems while your foot is malleable and able to adapt to a new position. Using the time while you are pregnant to support your foot in a new corrected position will allow proper functioning

and decreased pain for years to come.

A foot orthotic calibrated to your weight and foot flexibility is crucial to the foot during pregnancy and beyond. A premium custom orthotic will support your arch, effectively raising your arch. This raising will cause a shortening of the foot as it is placed in its optimal position and may actually reduce the need for bigger shoes. Improving your foot posture can also help unwanted postural changes in the knees and lower back.

Early preventative treatment with an orthotic is crucial for avoiding the development of other foot problems associated with over-pronation and arch drop such as plantar fasciitis (heel pain), bunion deformities and metatarsalgia (pain around the ball of the foot).



References

1. BIRD AR, MENZ HB, HYDE CC: The effect of Pregnancy on Footprint Parameters. J Amer Podiatr Med Assoc 89:405-409, 1999.
2. LYMBERY JK, GILLEARD W: The Stance Phase of Walking During Late Pregnancy. J Amer Podiatr Med Assoc 95(3): 247-253, 2005.
3. ANDERSON J DPM: www.babycenter.com "Ask the Experts" page.
4. FOTI T, DAVIDS JR, BAGLEY A: A Biomechanical Analysis of Gait During Pregnancy. J Bone and Joint Surg 82: 625-632, 2000.

This information page provides a general overview on this topic and may not apply to everyone. Always consult with your healthcare provider.